



## UWS Academic Portal

### **A trophic interaction framework for identifying the invasive capacity of novel organisms**

Penk, Marcin R. ; Saul, Wolf-Christian ; Dick, Jaimie T.A. ; Donohue, Ian ; Alexander, Mhairi E.; Linzmaier, Stefan ; Jeschke, Jonathan M.

*Published in:*  
Methods in Ecology and Evolution

*DOI:*  
[10.1111/2041-210X.12817](https://doi.org/10.1111/2041-210X.12817)

E-pub ahead of print: 20/05/2017

*Document Version*  
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

#### *Citation for published version (APA):*

Penk, M. R., Saul, W-C., Dick, J. T. A., Donohue, I., Alexander, M. E., Linzmaier, S., & Jeschke, J. M. (2017). A trophic interaction framework for identifying the invasive capacity of novel organisms. *Methods in Ecology and Evolution*. <https://doi.org/10.1111/2041-210X.12817>

#### **General rights**

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### **Take down policy**

If you believe that this document breaches copyright please contact [pure@uws.ac.uk](mailto:pure@uws.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# **A trophic interaction framework for identifying the invasive capacity of novel organisms**

Running title: An assessment framework for novel organisms

Word count: 6999

Marcin Penk<sup>\*1,2</sup>, Wolf-Christian Saul<sup>2,3,4</sup>, Jaimie T.A. Dick<sup>5</sup>, Ian Donohue<sup>1</sup>, Mhairi E. Alexander<sup>6</sup>, Stefan Linzmaier<sup>2,3,4</sup> and Jonathan M. Jeschke<sup>2,3,4</sup>

<sup>1</sup>School of Natural Sciences, Trinity College Dublin, College Green, Dublin 2, Ireland

<sup>2</sup>Department of Biology, Chemistry, Pharmacy, Institute of Biology, Freie Universität Berlin, Königin-Luise-Str. 1-3, 14195 Berlin, Germany

<sup>3</sup>Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Müggelseedamm 310, 12587 Berlin, Germany

<sup>4</sup>Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), Altensteinstr. 34, 14195 Berlin, Germany

<sup>5</sup>Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, MBC, 97 Lisburn Road, Belfast BT9 7BL, Northern Ireland, UK

<sup>6</sup>Institute for Biomedical and Environmental Health Research (IBEHR), School of Science and Sport, University of the West of Scotland, Paisley PA1 2BE, Scotland, UK

Corresponding author: penkm@tcd.ie

## Summary

1. The likelihood and impacts of invasions by novel organisms (e.g. non-native species, genetically-modified organisms) on the composition and functioning of receiving biological communities hinges on their capacity to exploit resources and/or avoid predation relative to resident counterparts. While assessment of invasion risk based on the comparison of functional responses (per-capita consumption rate as a function of resource density) of novel species with native analogues has been gaining popularity, it may be undermined if alternative prey and potential predators are not represented realistically.
2. Here, we propose a conceptual framework that enables rigorous identification of trophic traits conducive to invasion success by novel organisms – irrespective of their trophic position – and their likely ecological impacts, given their arrival and establishment. We focus on consumption here, but our framework can also be used for autotrophic energy acquisition, and extended to non-trophic and indirect interactions.
3. The framework enables a structured and prioritised selection of subsets of trophic links for invasion risk assessment. It is based on foraging theory and advances in comparative functional responses in invasion ecology. It can even be used in the absence of a resident comparator organism and when resources or predators are only partly known.
4. Our approach enhances the predictive power of species screening, and thus advances prevention and management of invasions under a common framework for all types of novel organisms.

**Key-words:** alien species, dietary generalism, ecological novelty, GMO, invasion success, functional responses, predator-prey trophic interactions, risk assessment

## Introduction

Predicting biological invasions (i.e. the spread of non-native species beyond the point of introduction) and managing their impacts (i.e. quantifiable alterations of the receiving ecosystem) remain key challenges in ecology (Simberloff *et al.* 2013). This demands improved understanding of the mechanisms of invasions. Human-assisted species translocations entail transfers across barriers that limit natural dispersal, and thus between environments which can have substantially different eco-evolutionary histories. Thus, introduced organisms can impart a high degree of ecological novelty to a system, which is conducive to invasiveness (Saul, Jeschke & Heger 2013). Organisms arriving in new environments enter resident ecological interaction networks, and the identification of their interactions within resident communities is important for understanding community dynamics. Predicting the attributes of these novel interactions is crucial for prioritising management of existing and anticipated invasions, and for assessing the side effects of intended introductions. Novel organisms (including translocated, but also range-expanding, genetically modified, synthesised or resurrected organisms; Jeschke, Keesing & Ostfeld 2013), whose ecological traits contrast with the eco-evolutionary experience of their resident interaction partners (Saul & Jeschke 2015), can potentially transform resident interaction networks through, for example, altering strengths, spatio-temporal patterns or other functional attributes of interactions (Mitchell *et al.* 2006; Downing *et al.* 2012; Mayer *et al.* 2013; Penk, Irvine & Donohue 2015).

Predation is a particularly important interaction type that can have strong impacts on community dynamics. This is primarily because it affects both predator fitness and prey biomass directly, potentially causing trophic cascades (Terborgh & Estes 2010; O'Connor *et al.* 2013) and food limitations for competitors (Strayer & Malcom 2007). Introduced predators can therefore affect resident prey populations significantly (Hays & Conant 2007;

Strayer 2009; Downing *et al.* 2012), with efficient exploitation of resources being conducive to high population growth, likely invasion success, and potentially considerable ecological impacts. Resource density is a key determinant of the feeding rate, and this relationship is characterised by ‘functional response’ curves (Holling 1959). For instance, decreasing prey density can reduce predator encounter rates with prey and thus offer a density-dependent prey refuge in a predator-prey system with a sigmoidal (i.e. Type III) functional response, but not with alternative functional response shapes where high proportions of prey are killed at low prey densities (Type I, II or their variants; Holling 1959; Jeschke, Kopp & Tollrian 2004; Jeschke & Tollrian 2005). Thus, the height – in particular the maximum feeding rate (i.e. curve asymptote) – and the shape of functional responses can reveal characteristics of consumer-resource interactions that are important for community dynamics and composition.

Invading predators with high ecological impact on their prey populations often have elevated functional responses compared to ecologically similar native species (Dick *et al.* 2017). Using comparative functional responses as an empirical screening method is thus rapidly gaining popularity among invasion biologists (Dick *et al.* 2013; Alexander *et al.* 2014; Barrios-O’Neill *et al.* 2014a; Rosewarne *et al.* 2016; Xu *et al.* 2016). The method typically infers invasion success and/or potential impacts from a limited number of prey species (frequently just one). However, biological invasions tend to lead to replacement of niche specialists by generalists (Clavel, Julliard & Devictor 2010), and numerous studies have identified a positive association between dietary generalism and invasion success (Bessa-Gomes *et al.* 2003; Jeschke & Strayer 2006; Romanuk *et al.* 2009; Clavel, Julliard & Devictor 2010; Arbaciauskas, Lesutiene & Gasiunaite 2013, but see Cassey *et al.* 2004 and Jackson *et al.* 2016). Thus, the inefficient use of one particular resident prey species does not necessarily preclude invasion success or impact upon ecological networks with realistic complexity. In other cases, a resident organism may be an inferior predator on a particular

prey species, and falsely appear a weaker overall interactor, compared to an introduced predator solely as a result of differing specialisation, despite apparent ecological similarity (Rosenfeld 2002; Dunoyer *et al.* 2014). Investigating functional responses with multiple prey, thereby taking into account the potential role of generalism and specialism, can buffer against such biases and at the same time improve the much needed representation of whole-ecosystem impacts of novel organisms (Ehrenfeld 2011; Simberloff 2011; Penk, Irvine & Donohue 2015). Furthermore, novel organisms can themselves be controlled by resident predators (Romanuk *et al.* 2009; MacNeil *et al.* 2013; Pintor & Byers 2015). Not accounting for top-down control experienced by introduced species, as has been typically the case in functional response-based screening methods, risks over-estimating their consumptive impacts and invasion success (but see Barrios-O'Neill *et al.* 2014b; Alexander, Raven & Robinson 2015).

Both top-down and bottom-up trophic interactions can thus directly affect the survival, fitness and ecological impacts of novel organisms. The complexity of these trophic links, including diet breadth and number of enemies, is an important determinant of invasion success (Romanuk *et al.* 2009). However, the logistics of incorporating multiple prey and predators into comparative functional responses may be demanding and frequently prohibitive.

We propose a conceptual framework for identifying and selecting a prioritised subset of trophic links to empirically identify the capacity for invasion success and ecological impacts of novel organisms (Fig. 1). We provide a worked example of the application of the framework for an intermediate consumer, the marbled crayfish (*Procambarus fallax* forma *virginalis*) in German low-land lakes (Fig. S1). This includes selection of both predators and prey, and can therefore instruct assessment of organisms of any trophic position. In addition, the marbled crayfish does not have any known native populations, and thus it is exemplary of

quite complex assessment scenarios. For clarity, our arguments relate to predation, which includes true predation, herbivory, parasitism and parasitoidism. However, detritivory can also be an important dietary subsidy of generalist consumers (Wise, Moldenhauer & Halaj 2006; Jackson *et al.* 2016) and should be carried through the assessment if it contributes to the diet of the novel ('focal') organism. Although we focus on consumption, our framework can be applied to autotrophic energy acquisition, and extended to non-trophic and indirect interactions. Whereas the non-empirical steps of our framework are readily applicable to any type of interaction, non-trophic interactions may require different empirical methods.

### **Mapping potential interaction partners in the target community**

Unless interaction with a particular resident organism is an *a priori* focus of assessment, an initial step of comprehensively mapping a potential network of direct consumptive interactions of the focal organism in the receiving ('target') community should enable minimisation of selection biases that may impede realistic assessment of the impact of the novel organism. This can be achieved by first listing all partners in direct consumptive interactions of the focal organism in its established range. This is then followed by matching all resident organisms in the target community that conform to the archetypes of these interaction partners and are likely to at least partly share spatio-temporal distribution patterns with the focal organism (Fig. 1). We define an archetype as organisms that have a similar set of morphological and behavioural traits that can condition a given type of interaction (Cox & Lima 2006; Winemiller *et al.* 2015), for example, feeding or defence strategy.

Observed trophic interactions in a given environment may not fully represent the feeding preferences of an organism (Futuyma & Moreno 1988; Devictor *et al.* 2010), and trophic interaction strength with a particular prey may depend on its availability in comparison to other prey rather than on the true preference of the consumer (Jaworski *et al.*

2013; Davis *et al.* 2015; Hanmer *et al.* 2017), as well as on environmental drivers. Thus, interaction partners of the focal organism as well as interaction strengths may vary among communities that differ in species composition and densities. If the focal organism is already established in the target environment, site-specific data about interaction partners should be given precedence above data from other areas. Otherwise, information from multiple communities within the distribution range of the focal species may improve control for context-dependencies. Assigning preference attributes based on how frequent and dominant interaction partners are throughout the established range (e.g. Kissling *et al.* 2014) can then help prioritise the selection of interaction partners for assessment. Considering ontogenetic stages of the focal organism with contrasting interaction partners (e.g. size class, identity or trophic guild of prey or enemies) could further improve predictions because limitation at any single stage preceding reproduction could constrict population dynamics (Werner & Gilliam 1984; Rudolf & Lafferty 2011).

For focal organisms that do not yet occur in nature, such as genetically modified, resurrected, synthetic, hybridised or selectively bred organisms, interaction partners of phylogenetically or functionally closest ('quasi-focal') organisms may provide reasonable approximation. For example, the marbled crayfish, introduced recently to German freshwaters (Chucholl, Morawetz & Groß 2012), originated in the aquarium trade and does not have any known native populations (Vogt *et al.* 2015). However, it is morphologically and functionally similar to the spiny-cheek crayfish (*Orconectes limosus*), a well-established earlier invader in Germany which can be considered a quasi-focal organism. The spiny-cheek crayfish is an omnivore that feeds on benthic invertebrates and macrophytes, and itself falls prey to fish, waterfowl and mustelids. Interaction partners of the spiny-cheek crayfish are thus good candidate prey and predators of the marbled crayfish (Fig. S1).



## Screening for a prioritised subset of the mapped interaction network

Should assessment be restricted to a subset of potential interaction partners, then criteria for selecting them depend on the goal of the assessment – invasion success, impact, or both (Fig. 1). The colonisation of areas beyond the point of introduction, synonymous with invasion success (Blackburn *et al.* 2011), is likely if an organism is able to utilise abundant resources, and/or if it can avoid high extrinsic mortality. Thus, we recommend focusing on potential resources with the highest biomass in the target ecosystem when selecting a prioritised subset of all identified potential interactions for the assessment of the likelihood of invasion. However, attention should also be paid to potential predators that are expected to have the highest predation pressure on the focal organism. Notably, high predation on the focal organism could come from consumers with high individual predation rates, or those that are not necessarily individually voracious but occur in high abundance (Dick *et al.* 2017).

Interactions of the focal organism with dominant predators and prey have the potential to affect major energy conduits within ecological networks and are thus conducive to strong ecosystem-level impacts, such as altered diversity, structure and functioning of target communities (Lockwood, Hoopes & Marchetti 2007; Penk, Irvine & Donohue 2015; Jackson *et al.* 2016). Further, interactions with keystone species or ecosystem engineers (Jones, Lawton & Shachak 1994; Power *et al.* 1996; Angelini *et al.* 2015), which are not necessarily very abundant, could amplify the indirect impact of invaders and convey ecosystem-level impacts. They should also be considered. It is also important to consider interactions with individual predator and prey organisms of conservation importance, regardless of whether such interactions have the potential to affect the whole community (Fig. 1).

In general, we advise selecting multiple prey and predator species of the focal organism for empirical assessment. However, their number and distribution among the interactor groups discussed above (i.e. those of highest biomass, keystone organisms and

ecosystem engineers, and those of conservation importance; Fig. 1), will depend on the purpose of the assessment, trophic level and niche breadth of the focal organism, food web complexity in the target ecosystem, management priorities and logistic constraints. For example, monophagous and oligophagous predators, including parasites and parasitoids, have inherently limited numbers of prey, while mesopredators typically have fewer predators than basal prey (Turney & Buddle 2016). Figure 2 shows exemplary hypothetical module structures for interaction settings between focal and resident organisms, indicating the diversity of interactions that need to be considered. Some of the interactor group categories will frequently overlap, and some may not be present in the target community. If the focal organism is already established, experimental trials or field data can be used to ascertain and prioritise interaction partners in the target community before engaging in full assessment.

Empirical examples of structured choices of prey in functional response studies of invasive species are rare (but see Dick *et al.* 2013; Barrios-O'Neill *et al.* 2016; Xu *et al.* 2016). In our worked example of the marbled crayfish (Fig. S1), the mussel *Dreissena* spp. has high abundance in the target community and is an ecosystem engineer. Dreissenids and other animal prey with poor escape response are key and preferred contributors to crayfish energy budgets (Momot 1995). The snails *Radix* spp. and *Bithynia tentaculata* are other important primary consumers in the target community that are readily consumed by crayfish (Olsen *et al.* 1991; Nyström, Brönmark & Granéli 1999). Testing predation on these three mollusc taxa could thus inform the assessment of both invasion success and ecological impacts of the marbled crayfish (Figs. S1 and S2). The quasi-focal organism (spiny-cheek crayfish) is a major prey of perch (*Perca fluviatilis*), which is a relatively abundant fish species in German low-land lakes and often holds key positions in food webs (Persson, Bystrom & Wahlstrom 2000). Predation by perch is likely restricted to immature or post-moult crayfish because of gape size limitation and the formidable defences of mature

crayfish. Nonetheless, this predatory fish could depress crayfish population dynamics, and thus it is a potentially important interactor (Figs. S1 and S2).

### **Empirically testing trophic interactions for the prioritised interaction subset**

Introduction of an organism that is of a predator or prey archetype already present in the resident community implies that resident prey or predators, respectively, are likely already familiar with such an archetype (Saul & Jeschke 2015). Because of such experience, it can be assumed that a novel organism can impact resident prey populations more strongly than their currently experienced predation pressure if its predatory traits toward a particular prey archetype are superior relative to its resident analogues. Similarly, a novel organism risks high impact from resident predators if its defences are weaker than those of its resident analogues, thus promoting prey switching. Therefore, comparing the trophic interaction strengths of the focal organism with its prey or predators, with those of an ecologically similar resident, where such exists, provides a useful benchmark for gauging the magnitude of interaction strength (van Kleunen *et al.* 2010; Dick *et al.* 2014). By definition, no two species are identical (Ordóñez 2014), but resident organisms that are of the same predator or prey archetype (Cox & Lima 2006; Winemiller *et al.* 2015) can offer a useful approximation of a reference baseline if any relevant functional differences between otherwise analogous species are acknowledged. In our worked example, the marbled crayfish co-occurs with other omnivorous crayfish (Chucholl, Morawetz & Groß 2012) of a similar predator and prey archetype that can be used as comparators (Fig. S1).

On the other hand, a novel organism that does not have any resident comparator is likely to have characteristics largely unfamiliar to resident prey and predators and thus the potential to bypass their defences and offences (Saul & Jeschke 2015). In such a case, the absolute, rather than comparative interaction strength of the focal novel organism with its

prey and predators can be of primary interest, and offtake rate of prey in relation to its reproductive rate can be used to predict impact on prey populations (MacNeil *et al.* 2013; Fig. 1). Qualitative pilot experiments can inform which degree of functional similarity can be assumed as a baseline.

## **Inference to real ecosystems**

*In situ* measurements and manipulations provide realistic settings, but tend to allow poor control of confounding factors (but see Barrios-O'Neill *et al.* 2014a). Also, they cannot be carried out if the focal organism is not (yet) present in the target environment. Laboratory experiments, on the other hand, typically simplify biotic and abiotic contexts, and the applied relevance of their results depends on the degree to which experimental settings facilitate natural offensive and defensive behaviour. For example, sheltering or camouflage may alter the shape of density-dependent predation, in that individual organisms devoid of their typical protective settings during experiments are more exposed to predation (Whittingham & Markland 2002; Horppila *et al.* 2003; Alexander, Dick & O'Connor 2013; Barrios-O'Neill *et al.* 2015). Both ambient temperatures and environmental hypoxia can also affect activity level, and moderate predator-prey interactions (Englund *et al.* 2011; Lavery *et al.* 2015; Penk *et al.* 2016). Laboratory-derived functional responses typically isolate an individual predator and single prey species (e.g. Dick *et al.* 2013; Barrios-O'Neill *et al.* 2014a; Xu *et al.* 2016, but see Alexander, Dick & O'Connor 2013; Medoc, Spataro & Arditi 2013; Wasserman *et al.* 2016), and thus rarely account for prey switching or interference among predators which could affect the outcome of an interaction (Amarasekare 2002; Tschanz, Bersier & Bacher 2007; van Leeuwen *et al.* 2013). The degree of spatio-temporal overlap of habitat use by the focal organism and its interaction partners is another important consideration (Polis, Anderson & Holt 1997). For example, a potentially strong interactor may have only a small

time window for realising such interactions if it rarely encounters particular prey and predator species. Whereas detailed propositions for ameliorating these problems are beyond the scope of this manuscript, we emphasise that lack of their consideration can undermine inference.

On the other hand, the need to quantify absolute interaction strengths accurately is circumvented in comparative studies, which focus on consumption rates relative to a native analogue rather than on absolute values, with an underlying assumption that both comparators would be influenced similarly by experimental artefacts (Dick *et al.* 2014). Indeed, comparative functional responses derived from simple laboratory experiments can be highly successful in explaining real-ecosystem ecological impacts of invaders across taxonomic and trophic groups (Dick *et al.* 2017). Context-dependencies may thus be particularly influential in making inference from studies on a novel organism that does not have a resident comparator because they rely on quantification of absolute interaction strengths.

Our framework specifically focuses on biological interactions, but the importance of intrinsic characteristics of the novel organism also has to be considered. For example, prognoses of population and community dynamics require at least some information on the reproductive rates of the focal organism and its interaction partners. The reproductive rate of a consumer determines the degree to which it can capitalise numerically on its ability to exploit prey and cumulatively increase its impact on prey populations, whereas the reproductive rate of prey determines their capacity to persist under given predation pressure (Twardochleb, Novak & Moore 2012). Both of these factors are key drivers of community dynamics.

Any model necessitates a trade-off between generality, realism and precision (Levins 1966). It is impossible to achieve all of these simultaneously to full extent, and the decision as to how to optimise this trade-off depends upon the focal system. We therefore present a

basic framework here, which needs to be adjusted and extended on a case-by-case basis to make it useful for the particular focal system in question. For example, for many systems it will be useful to incorporate non-consumptive or indirect interactions, or impacts on ecosystem services into the basic framework.

### **Non-consumptive and indirect interactions**

Consumptive interactions are the key focus of our framework (Fig. 1). However, non-consumptive and indirect (trait-mediated) interactions, for example through interference, facilitation and inhibition, can have important implications for community dynamics and in some cases even take prominence over direct impacts (Suraci *et al.* 2016). Indirect interactions occur when one species alters the effects that another species has on a third, potentially confounding predicted impacts of a novel organism that are derived from two-species studies (White, Wilson & Clarke 2006). For instance, changes to the foraging behaviour of a resident intermediate consumer as a result of the presence of a novel higher-order predator may alter the strength of interactions with a basal prey resource, releasing it from predation pressure (Townsend 1996). Alternatively, the presence of a resident higher-order predator may result in an exacerbated effect of a non-resident intermediate species towards its prey in comparison to a resident consumer, again influencing impact of the focal organism (Barrios-O'Neill *et al.* 2014b). Quantification of beneficial and disadvantageous outcomes of such interactions, in particular regulation of feeding and mortality rates, could be readily integrated in the empirical steps of our framework.

### **Conclusions**

Key theoretical progress on functional responses in invasion ecology has come from retrospective empirical attempts to explain invasion success and impacts of established

invaders (Hooff & Bollens 2004; Radford, Dickinson & Lord 2007; Bollache *et al.* 2008; Dick *et al.* 2013). Such attempts typically focus on isolated interactions in which the invader is clearly efficient and superior over a native comparator. However, biological invasions are highly dependent on biological contexts (Donohue *et al.* 2013; Ricciardi *et al.* 2013; Saul, Jeschke & Heger 2013), and robust prospective applications require a more comprehensive assessment network with multiple interaction partners, including predators of the focal organism. Applications of functional responses in biocontrol frequently fail to explain impact on individual prey organisms (Lester & Harmsen 2002; Fernández-Arhex & Corley 2003). The inclusion of predators and alternative prey, together with more realistic representation of key abiotic conditions and explicit discussion of the relevance of results to natural ecosystems can improve explanatory and predictive power of impact assessments. We focused here on predation in a broad sense, but the same assessment protocol and analogous empirical methods can be used for detritivorous and autotrophic energy acquisition (Radford, Dickinson & Lord 2007; McNickle & Brown 2014).

Risk assessment based on performance in comparable environments, where such information exists, is less laborious than collecting new data. However, interaction partners in new and existing ranges should be compared in a structured way to minimise bias. The steps of our framework that are based on existing data can be used to inform such comparisons (Fig. 1). Furthermore, novel organisms can be introduced to dissimilar communities or abiotic conditions in comparison to their existing ranges, or they can be absent in nature. Such scenarios preclude comparisons based on performance elsewhere and necessitate collection of new data (Fig. 1). In the face of limiting resources, a compromise between experimental complexity and accuracy of risk assessments needs to be reached on a case-by-case basis. Notably, relevant empirical data can be collected *in situ* (Angerbjorn, Tannerfeldt & Erlinge 1999, Goss-Custard *et al.* 2006; Moustahfid *et al.* 2010, Barrios-O'Neill *et al.*

2014a), permitting empirical testing of organisms that do not lend themselves well to laboratory conditions, or should not be interfered with on ethical grounds. In any case, the broader interaction network in the focal ecosystem should be at least theoretically considered, even if just to critically scrutinise the assessment outcomes. The non-empirical steps of our framework can inform such exercises regardless of the scale or complexity of the system in question.

## **Acknowledgments**

We appreciate helpful comments by the Associate Editor and two reviewers. MP was supported by a COST Action TD1209 STSM grant (011015-062070). WCS and JMJ were supported by the ERA-Net BiodivERsA (project FFII) with Deutsche Forschungsgemeinschaft (DFG; JE 288/7-1). JMJ was additionally supported by the DFG grant JE 288/9-1.

## **Data Accessibility**

This manuscript does not include any data.

## **Author contributions statement**

MP, WCS and JMJ conceived key ideas; MP led the writing of the manuscript; all authors contributed critically to the drafts.

## **References**

Alexander, M.E., Dick, J.T.A. & O'Connor, N.E. (2013) Trait-mediated indirect interactions in a marine intertidal system as quantified by functional responses. *Oikos*, **122**, 1521–1531.



371 Alexander, M.E., Dick, J.T.A., Weyl, O.L.F., Robinson, T.B. & Richardson, D.M. (2014)  
 372 Existing and emerging high impact invasive species are characterized by higher functional  
 373 responses than natives. *Biology Letters*, **10**, doi: 10.1098/rsbl.2013.0946.  
 374  
 375 Alexander, M.E., Raven, H.J. & Robinson, T.B. (2015) Foraging decisions of a native whelk,  
 376 *Trochia cingulata* Linnaeus, and the effects of invasive mussels on prey choice. *Journal of*  
 377 *Experimental Marine Biology and Ecology*, **470**, 26–33.  
 378  
 379 Amarasekare, P. (2002) Interference competition and species coexistence. *Proceedings of the*  
 380 *Royal Society B: Biological Sciences*, **269**, 2541–2550.  
 381  
 382 Angelini, C., van der Heide, T., Griffin, J.N., Morton, J.P., Derksen-Hooijberg, M., Lamers,  
 383 L.P.M., Smolders, A.J.P. & Silliman, B.R. (2015) Foundation species' overlap enhances  
 384 biodiversity and multifunctionality from the patch to landscape scale in southeastern United  
 385 States salt marshes. *Proceedings of the Royal Society of London B: Biological Sciences*, **282**,  
 386 doi: 10.1098/rspb.2015.0421.  
 387  
 388 Angerbjorn, A., Tannerfeldt, M. & Erlinge, S. (1999) Predator-prey relationships: Arctic  
 389 foxes and lemmings. *Journal of Animal Ecology*, **68**, 34–49.  
 390  
 391 Arbaciauskas, K., Lesutiene, J. & Gasiunaite, Z.R. (2013) Feeding strategies and elemental  
 392 composition in Ponto-Caspian peracaridans from contrasting environments: can  
 393 stoichiometric plasticity promote invasion success? *Freshwater Biology*, **58**, 1052–1068.  
 394  
 395 Barrios-O'Neill, D., Dick, J.T.A., Ricciardi, A., MacIsaac, H.J. & Emmerson, M.C. (2014a)

396 Deep impact: in situ functional responses reveal context-dependent interactions between  
 397 vertically migrating invasive and native mesopredators and shared prey. *Freshwater Biology*,  
 398 **59**, 2194–2203.

399

400 Barrios-O'Neill, D., Dick, J.T.A., Emmerson, M.C., Ricciardi, A., MacIsaac, H.J., Alexander,  
 401 M.E. & Bovy, H.C. (2014b) Fortune favours the bold: a higher predator reduces the impact of  
 402 a native but not an invasive intermediate predator. *Journal of Animal Ecology*, **83**, 693–701.

403

404 Barrios-O'Neill, D., Dick, J.T.A., Emmerson, M.C., Ricciardi, A. & MacIsaac, H.J. (2015)  
 405 Predator-free space, functional responses and biological invasions. *Functional Ecology*, **29**,  
 406 377–384.

407

408 Barrios-O'Neill, D., Kelly, R., Dick, J.T.A., Ricciardi, A., MacIsaac, H.J. & Emmerson, M.C.  
 409 (2016) On the context-dependent scaling of consumer feeding rates. *Ecology Letters*, **19**,  
 410 668–678.

411

412 Bessa-Gomes, C., Danek-Gontard, M., Cassey, P., Moller, A.P., Legendre, S. & Clobert, J.  
 413 (2003) Mating behaviour influences extinction risk: insights from demographic modelling  
 414 and comparative analysis of avian extinction risk. *Annales Zoologici Fennici*, **40**, 231–245.

415

416 Blackburn, T.M., Pyšek, P., Bacher, S., Carlton, J.T., Duncan, R.P., Jarošík, V., Wilson,  
 417 J.R.U. & Richardson, D.M. (2011) A proposed unified framework for biological invasions.  
 418 *Trends in Ecology and Evolution*, **26**, 333–339.

419

420 Bollache, L., Dick, J.T.A., Farnsworth, K.D. & Montgomery, W.I. (2008) Comparison of the

functional responses of invasive and native amphipods. *Biology Letters*, **4**, 166–169.

Cassey, P., Blackburn, T.M., Sol, D., Duncan, R.P. & Lockwood, J.L. (2004) Global patterns of introduction effort and establishment success in birds. *Proceedings of the Royal Society B: Biological Sciences*, **271**, S405–S408.

Chucholl, C., Morawetz, K. & Groß, H. (2012) The clones are coming – strong increase in marmorkrebs [*Procambarus fallax* (Hagen, 1870) f. *virginalis*] records from Europe. *Aquatic Invasions*, **7**, 511–519.

Clavel, J., Julliard, R. & Devictor, V. (2010) Worldwide decline of specialist species: toward a global functional homogenization? *Frontiers in Ecology and the Environment*, **9**, 222–228.

Cox, J.G. & Lima, S.L. (2006) Naiveté and an aquatic-terrestrial dichotomy in the effects of introduced predators. *Trends in Ecology and Evolution*, **21**, 674–680.

Davis, N.E., Forsyth, D.M., Triggs, B., Pascoe, C., Benshemesh, J., Robley, A., Lawrence, J., Ritchie, E.G., Nimmo, D.G. & Lumsden, L.F. (2015) Interspecific and geographic variation in the diets of sympatric carnivores: dingoes/wild dogs and red foxes in south-eastern Australia. *PLoS ONE*, **10**, e0120975.

Devictor, V., Clavel, J., Julliard, R., Lavergne, S., Mouillot, D., Thuiller, W., Venail, P., Villéger, S. & Mouquet, N. (2010) Defining and measuring ecological specialization. *Journal of Applied Ecology*, **47**, 15–25.

446 Dick, J.T.A., Gallagher, K., Avlijas, S., Clarke, H.C., Lewis, S.E., Leung, S., Minchin, D.,  
 447 Caffrey, J., Alexander, M.E., Maguire, C., Harrod, C., Reid, N., Haddaway, N.R.,  
 448 Farnsworth, K.D., Penk, M. & Ricciardi, A. (2013) Ecological impacts of an invasive  
 449 predator explained and predicted by comparative functional responses. *Biological Invasions*,  
 450 **15**, 837–846.  
 451  
 452 Dick, J.T.A., Alexander, M.E., Jeschke, J.M., Ricciardi, A., MacIsaac, H.J., Robinson, T.B.,  
 453 Kumschick, S., Weyl, O.L.F., Dunn, A.M., Hatcher, M.J., Paterson, R.A., Farnsworth, K.D.  
 454 & Richardson, D.M. (2014) Advancing impact prediction and hypothesis testing in invasion  
 455 ecology using a comparative functional response approach. *Biological Invasions*, **16**, 735–  
 456 753.  
 457  
 458 Dick, J.T.A., Lavery, C., Lennon, J.J., Barrios-O'Neill, D., Mensink, P.J., Britton, J.R.,  
 459 Medoc, V., Boets, P., Alexander, M.E., Taylor, N.G., Dunn, A.M., Hatcher, M.J., Rosewarne,  
 460 P.J., Crookes, S., MacIsaac, H.J., Xu, M., Ricciardi, A., Wasserman, R.J., Ellender, B.R.,  
 461 Weyl, O.L.F., Lucy, F.E., Banks, P.B., Dodd, J.A., MacNeil, C., Penk, M.R., Aldridge, D.C.  
 462 & Caffrey, J.M. (2017). Invader Relative Impact Potential: a new metric to understand and  
 463 predict the ecological impacts of existing, emerging and future invasive alien species. *Journal*  
 464 *of Applied Ecology*, doi: 10.1111/1365-2664.12849.  
 465  
 466 Donohue, I., Petchey, O.L., Montoya, J.M., Jackson, A.L., McNally, L., Viana, M., Healy,  
 467 K., Lurgi, M., O'Connor, N.E. & Emmerson, M.C. (2013) On the dimensionality of  
 468 ecological stability. *Ecology Letters*, **16**, 421–429.  
 469  
 470 Downing, A.S., van Nes, E.H., Janse, J.H., Witte, F., Cornelissen, I.J.M., Scheffer, M. &

471 Mooij, W.M. (2012) Collapse and reorganization of a food web of Mwanza Gulf, Lake  
 472 Victoria. *Ecological Applications*, **22**, 229–239.

473

474 Dunoyer, L., Dijoux, L., Bollache, L. & Lagrue, C. (2013) Effects of crayfish on leaf litter  
 475 breakdown and shredder prey: are native and introduced species functionally redundant?  
 476 *Biological Invasions*, **16**, 1545–1555.

477

478 Ehrenfeld, J.G. (2011) Ecosystem consequences of biological invasions. *Annual Reviews of*  
 479 *Ecology, Evolution, and Systematics*, **41**, 59–80.

480

481 Englund, G., Öhlund, G., Hein, C.L. & Diehl, S. (2011) Temperature dependence of the  
 482 functional response. *Ecology Letters*, **14**, 914–921.

483

484 Fernández-Arhex, V. & Corley, J.C. (2003) The functional response of parasitoids and its  
 485 implications for biological control. *Biocontrol Science and Technology*, **13**, 403–413.

486

487 Futuyma, D.J. & Moreno, G. (1988) The evolution of ecological specialization. *Annual*  
 488 *Review of Ecology and Systematics*, **19**, 207–233.

489

490 Goss-Custard, J.D., West, A.D., Yates, M.G., Caldow, R.W., Stillman, R.A., Bardsley, L.,  
 491 Castilla, J., Castro, M., Dierschke, V., Durell, S.E., Eichhorn, G., Ens, B.J., Exo, K.M.,  
 492 Udayangani-Fernando, P., Ferns, P.N., Hockey, P.A., Gill, J.A., Johnstone, I., Kalejta-  
 493 Summers, B., Masero, J.A., Moreira, F., Nagarajan, R.V., Owens, I.P., Pacheco, C., Perez-  
 494 Hurtado, A., Rogers, D., Scheiffarth, G., Sitters, H., Sutherland, W.J., Triplet, P., Worrall,  
 495 D.H., Zharikov, Y., Zwarts, L. & Pettifor, R.A. (2006) Intake rates and the functional

496 response in shorebirds (Charadriiformes) eating macro-invertebrates. *Biological Reviews*, **81**,  
 497 501–529.  
 498  
 499 Hanmer, J., White, J.W. & Pawlik, J.R. (2017) Application of diet theory reveals context-  
 500 dependent foraging preferences in an herbivorous coral reef fish. *Oecologia*, **184**, 127–137.  
 501  
 502 Hays, W.S.T. & Conant, S. (2007) Biology and impacts of pacific island invasive species: 1.  
 503 a worldwide review of effects of the small Indian mongoose, *Herpestes javanicus* (Carnivora:  
 504 Herpestidae). *Pacific Science*, **61**, 3–16.  
 505  
 506 Holling, C.S. (1959) The components of predation as revealed by a study of small-mammal  
 507 predation of the European pine sawfly. *The Canadian Entomologist*, **91**, 293–320.  
 508  
 509 Hooff, R., C. & Bollens, S., M. (2004) Functional response and potential predatory impact of  
 510 *Tortanus dextrilobatus*, a carnivorous copepod recently introduced to the San Francisco  
 511 Estuary. *Marine Ecology Progress Series*, **277**, 167–179.  
 512  
 513 Horppila, J., Liljendahl-Nurminen, A., Malinen, T., Salonen, M., Tuomaala, A., Uusitalo, L.  
 514 & Vinni, M. (2003) *Mysis relicta* in a eutrophic lake: consequences of obligatory habitat  
 515 shifts. *Limnology and Oceanography*, **48**, 1214–1222.  
 516  
 517 Jackson, M.C., Grey, J., Miller, K., Britton, J.R. & Donohue, I. (2016) Dietary niche  
 518 constriction when invaders meet natives: evidence from freshwater decapods. *Journal of*  
 519 *Animal Ecology*, **85**, 1098–1107.  
 520

521 Jaworski, C.C., Bompard, A., Genies, L., Amiens-Desneux, E. & Desneux, N. (2013)  
522 Preference and prey switching in a generalist predator attacking local and invasive alien  
523 pests. *PLoS ONE*, **8**, e82231.

524

525 Jeschke, J.M., Kopp, M. & Tollrian, R. (2004) Consumer-food systems: why type I  
526 functional responses are exclusive to filter feeders. *Biological Reviews*, **79**, 337–349.

527

528 Jeschke, J.M & Tollrian, R. (2005) Effects of predator confusion on functional responses.  
529 *Oikos*, **111**, 547–555.

530

531 Jeschke, J.M. & Strayer, D.L. (2006) Determinants of vertebrate invasion success in Europe  
532 and North America. *Global Change Biology*, **12**, 1608–1619.

533

534 Jeschke, J.M., Keesing, F. & Ostfeld, R.S. (2013) Novel organisms: comparing invasive  
535 species, GMOs, and emerging pathogens. *Ambio*, **42**, 541–548.

536

537 Jones, C.G., Lawton, J.H. & Shachak, M. (1994) Organisms as ecosystem engineers. *Oikos*,  
538 **69**, 373–386.

539

540 Kissling, W.D., Dalby, L., Fløjgaard, C., Lenoir, J., Sandel, B., Sandom, C., Trøjelsgaard, K.  
541 & Svenning, J.-C. (2014) Establishing macroecological trait datasets: digitalization,  
542 extrapolation, and validation of diet preferences in terrestrial mammals worldwide. *Ecology*  
543 *and Evolution*, **4**, 2913–2930.

544

545 Laverty, C., Dick, J.T.A., Alexander, M.E. & Lucy, F.E. (2015) Differential ecological

546 impacts of invader and native predatory freshwater amphipods under environmental change  
547 are revealed by comparative functional responses. *Biological Invasions*, **17**, 1761–1770.  
548

549 Lester, P.J. & Harmsen, R. (2002) Functional and numerical responses do not always indicate  
550 the most effective predator for biological control: an analysis of two predators in a two-prey  
551 system. *Journal of Applied Ecology*, **39**, 455–468.  
552

553 Levins, R. (1966) The strategy of model building in population biology. *American Scientist*,  
554 **54**, 421–431.  
555

556 Lockwood, J.L., Hoopes, M.F. & Marchetti, M.P. (2007) *Invasion Ecology*. Blackwell  
557 Publishing, Oxford.  
558

559 MacNeil, C., Dick, J.T.A., Alexander, M., Dodd, J. & Ricciardi, A. (2013) Predators vs.  
560 alien: differential biotic resistance to an invasive species by two resident predators. *NeoBiota*,  
561 **19**, 1–19.  
562

563 Mayer, C., M., Burlakova, L., E., Eklöv, P., Fitzgerald, D., Karatayev, A., Y., Ludsine, S., A.,  
564 Millard, S., Mills, E., L., Ostapenko, A.P., Rudstam, L., G., Zhu, B. & Zhukova, T., V.  
565 (2013) Benthification of freshwater lakes. *Quagga and Zebra Mussels: Biology, Impacts, and*  
566 *Control* (eds T.F. Nalepa & D.W. Schloesser), pp. 575–586. CRC Press, Boca Raton, FL.  
567

568 McNickle, G.G. & Brown, J.S. (2014) When Michaelis and Menten met Holling: towards a  
569 mechanistic theory of plant nutrient foraging behaviour. *AoB PLANTS*, **6**, doi:  
570 10.1093/aobpla/plu066.



571

572 Medoc, V., Spataro, T. & Arditi, R. (2013) Prey: predator ratio dependence in the functional  
 573 response of a freshwater amphipod. *Freshwater Biology*, **58**, 858–865.

574

575 Mitchell, C.E., Agrawal, A.A., Bever, J.D., Gilbert, G.S., Hufbauer, R.A., Klironomos, J.N.,  
 576 Maron, J.L., Morris, W.F., Parker, I.M., Power, A.G., Seabloom, E.W., Torchin, M.E. &  
 577 Vázquez, D.P. (2006) Biotic interactions and plant invasions. *Ecology Letters*, **9**, 726–740.

578

579 Momot, W.T. (1995) Redefining the role of crayfish in aquatic ecosystems. *Reviews in*  
 580 *Fisheries Science*, **3**, 33–63.

581

582 Moustahfid, H., Tyrrell, M.C., Link, J.S., Nye, J.A., Smith, B.E. & Gamble, R.J. (2010)  
 583 Functional feeding responses of piscivorous fishes from the northeast US continental shelf.  
 584 *Oecologia*, **163**, 1059–1067.

585

586 Nyström, P., Brönmark, C. & Granéli, W. (1999) Influence of an exotic and a native crayfish  
 587 species on a littoral benthic community. *Oikos*, **85**, 545–553.

588

589 O'Connor, N.E., Emmerson, M.C., Crowe, T.P. & Donohue, I. (2013) Distinguishing  
 590 between direct and indirect effects of predators in complex ecosystems. *Journal of Animal*  
 591 *Ecology*, **82**, 438–448.

592

593 Olsen, T.M., Lodge, D.M., Capelli, G.M. & Houlihan, R.J. (1991) Mechanisms of impact of  
 594 an introduced crayfish (*Orconectes rusticus*) on littoral congeners, snails, and macrophytes.  
 595 *Canadian Journal of Fisheries and Aquatic Sciences*, **48**, 1853–1861.

596

597   Ordóñez, A. (2014) Functional and phylogenetic similarity of alien plants to co-occurring  
598   natives. *Ecology*, **95**, 1191–1202.

599

600   Penk, M., Irvine, K. & Donohue, I. (2015) Ecosystem-level effects of a globally spreading  
601   invertebrate invader are not moderated by a functionally similar native. *Journal of Animal*  
602   *Ecology*, **84**, 1628–1636.

603

604   Penk, M.R., Jeschke, J.M., Minchin, D. & Donohue, I. (2016) Warming can enhance invasion  
605   success through asymmetries in energetic performance. *Journal of Animal Ecology*, **85**, 419–  
606   426.

607

608   Persson, L., Bystrom, P. & Wahlstrom, E. (2000) Cannibalism and competition in Eurasian  
609   perch: population dynamics of an ontogenetic omnivore. *Ecology*, **81**, 1058–1071.

610

611   Pintor, L.M. & Byers, J.E. (2015) Do native predators benefit from non-native prey? *Ecology*  
612   *Letters*. **18**, 1174–1180.

613

614   Polis, G.A., Anderson, W.B. & Holt, R.D. (1997) Towards an integration of landscape and  
615   food web ecology: the dynamics of spatially subsidized food webs. *Annual Review of*  
616   *Ecology and Systematics*, **28**, 289–316.

617

618   Power, M.E., Tilman, D., Estes, J.A., Menge, B.A., Bond, W.J., Mills, L.S., Daily, G.,  
619   Castilla, J.C., Lubchenco, J. & Paine, R.T. (1996) Challenges in the quest for keystones:  
620   identifying keystone species is difficult—but essential to understanding how loss of species  
621   will affect ecosystems. *Bioscience*, **46**, 609–620.

622

623 Radford, I.J., Dickinson, K.J.M. & Lord, J.M. (2007) Functional and performance  
 624 comparisons of invasive *Hieracium lepidulum* and co-occurring species in New Zealand.  
 625 *Austral Ecology*, **32**, 338–354.

626

627 Ricciardi, A., Hoopes, M.F., Marchetti, M.P. & Lockwood, J.L. (2013) Progress toward  
 628 understanding the ecological impacts of nonnative species. *Ecological Monographs*, **83**, 263–  
 629 282.

630

631 Romanuk, T.N., Zhou, Y., Brose, U., Berlow, E.L., Williams, R.J. & Martinez, N.D. (2009)  
 632 Predicting invasion success in complex ecological networks. *Philosophical Transactions of*  
 633 *the Royal Society of London B: Biological Sciences*, **364**, 1743–1754.

634

635 Rosenfeld, J.S. (2002) Functional redundancy in ecology and conservation. *Oikos*, **98**, 156–  
 636 162.

637

638 Rosewarne, P.J., Mortimer, R.J.G., Newton, R.J., Grocock, C., Wing, C.D. & Dunn, A.M.  
 639 (2016) Feeding behaviour, predatory functional responses and trophic interactions of the  
 640 invasive Chinese mitten crab (*Eriocheir sinensis*) and signal crayfish (*Pacifastacus*  
 641 *leniusculus*). *Freshwater Biology*, **61**, 426–443.

642

643 Rudolf, V.H.W. & Lafferty, K.D. (2011) Stage structure alters how complexity affects  
 644 stability of ecological networks. *Ecology Letters*, **14**, 75–79.

645

646 Saul, W.-C., Jeschke, J.M. & Heger, T. (2013) The role of eco-evolutionary experience

647 in invasion success. *NeoBiota*, **17**, 57–74.

648

649 Saul, W.-C. & Jeschke, J.M. (2015) Eco-evolutionary experience in novel species

650 interactions. *Ecology Letters*, **18**, 236–245.

651

652 Simberloff, D. (2011) How common are invasion-induced ecosystem impacts? *Biological*

653 *Invasions*, **13**, 1255–1268.

654

655 Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J.,

656 Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E.

657 & Vilà, M. (2013) Impacts of biological invasions: what's what and the way forward. *Trends*

658 *in Ecology and Evolution*, **28**, 58–66.

659

660 Strayer, D.L. & Malcom, H.M. (2007) Effects of zebra mussels (*Dreissena polymorpha*) on

661 native bivalves: the beginning of the end or the end of the beginning? *Journal of the North*

662 *American Benthological Society*, **26**, 111–122.

663

664 Strayer, D.L. (2009) Twenty years of zebra mussels: lessons from the mollusk that made

665 headlines. *Frontiers in Ecology and the Environment*, **7**, 135–141.

666

667 Suraci, J.P., Clinchy, M., Dill, L.M., Roberts, D. & Zanette, L.Y. (2016) Fear of large

668 carnivores causes a trophic cascade. *Nature Communications*, **7**, 10698.

669

670 Terborgh, J. & Estes, J.A. (2010) *Trophic Cascades: Predators, Prey, and the Changing*

671 *Dynamics of Nature*. Island Press, Washington DC.

672

673 Townsend, C.R. (1996) Invasion biology and ecological impacts of brown trout *Salmo trutta*

674 in New Zealand. *Biological Conservation*, **78**, 13–22.

675

676 Tschanz, B., Bersier, L.-F. & Bacher, S. (2007) Functional responses: a question of

677 alternative prey and predator density. *Ecology*, **88**, 1300–1308.

678

679 Turney, S. & Buddle, C.M. (2016) Pyramids of species richness: the determinants and

680 distribution of species diversity across trophic levels. *Oikos*, **125**, 1224–1232.

681

682 Twardochleb, L.A., Novak, M. & Moore, J.W. (2012) Using the functional response of a

683 consumer to predict biotic resistance to invasive prey. *Ecological Applications*, **22**, 1162–

684 1171.

685

686 van Kleunen, M., Dawson, W., Schlaepfer, D., Jeschke, J.M. & Fischer, M. (2010) Are

687 invaders different? A conceptual framework of comparative approaches for assessing

688 determinants of invasiveness. *Ecology Letters*, **13**, 947–958.

689

690 van Leeuwen, E., Brännström, Å., Jansen, V.A.A., Dieckmann, U. & Rossberg, A.G. (2013)

691 A generalized functional response for predators that switch between multiple prey species.

692 *Journal of Theoretical Biology*, **328**, 89–98.

693

694 Vogt, G., Falckenhayn, C., Schrimpf, A., Schmid, K., Hanna, K., Panteleit, J., Helm, M.,

695 Schulz, R. & Lyko, F. (2015) The marbled crayfish as a paradigm for saltational speciation

696 by autopolyploidy and parthenogenesis in animals. *Biology Open*, **4**, 1583–1594.

697

698 Wasserman, R.J., Alexander, M.E., Dalu, T., Ellender, B.R., Kaiser, H. & Weyl, O.L.F.

699 (2016) Using functional responses to quantify interaction effects among predators.

700 *Functional Ecology*, doi: 10.1111/1365-2435.12682.

701

702 Werner, E.E. & Gilliam, J.F. (1984) The ontogenetic niche and species interactions in size-

703 structured populations. *Annual Review of Ecology and Systematics*, **15**, 393–425.

704

705 White, E.M., Wilson, J.C. & Clarke, A.R. (2006) Biotic indirect effect: a neglected concept in

706 invasion biology. *Diversity and Distributions*, **12**, 443–445.

707

708 Whittingham, M. & Markland, H. (2002) The influence of substrate on the functional

709 response of an avian granivore and its implications for farmland bird conservation.

710 *Oecologia*, **130**, 637–644.

711

712 Winemiller, K.O., Fitzgerald, D.B., Bower, L.M. & Pianka, E.R. (2015) Functional traits,

713 convergent evolution, and periodic tables of niches. *Ecology Letters*, **18**, 737–751.

714

715 Wise, D.H., Moldenhauer, D.M. & Halaj, J. (2006) Using stable isotopes to reveal shifts in

716 prey consumption by generalist predators. *Ecological Applications*, **16**, 865–876.

717

718 Xu, M., Mu, X., Dick, J.T.A., Fang, M., Gu, D., Luo, D., Zhang, J., Luo, J. & Hu, Y. (2016)

719 Comparative functional responses predict the invasiveness and ecological impacts of alien

720 herbivorous snails. *PLoS ONE*, **11**, e0147017.

721

## Supporting Information

**Fig. S1.** Illustration of predator and prey selection using an example of the marbled crayfish.

**Fig. S2.** Illustration of an interaction module structure for testing, using an example of the marbled crayfish.

## Figure captions

**Fig. 1.** Framework for quantifying interaction strength of a focal organism with multiple resident organisms in the target community for a given type of interaction. Nodes and broken links indicate alternative and supplementary paths, respectively. *S* and *I* indicate interaction-partner categories relevant for the assessment of invasion success and ecological impacts, respectively.

**Fig. 2.** Hypothetical interaction module structures for testing trophic interactions of a focal organism (oval shapes) representing top (a, b), intermediate (c, d) and basal (e) trophic positions, and two extrema on the diet-breadth continuum (polyphagous [a, c] and monophagous [b, d]).




Fig. 1






Fig. 2

774 **Supporting Information**

775

776 A trophic interaction framework for identifying the invasive capacity of novel organisms

777

778 Marcin Penk, Wolf-Christian Saul, Jaimie T.A. Dick, Ian Donohue, Mhairi E. Alexander,

779 Stefan Linzmaier and Jonathan M. Jeschke




Fig. S1. Illustration of the framework for quantifying interaction strength of a focal organism with resident organisms in the target community, using an example of predator and prey selection for the marbled crayfish (*Procambarus fallax* forma *virginalis*) in a German low-land lake. *S* and *I* indicate prey categories relevant for the assessment of invasion success and ecological impacts, respectively. Quasi-focal organism is also the resident comparator in this example.




Fig. S2. Illustration of the interaction module structure for testing using an example of predator and prey (top and bottom boxes, respectively) selected for the marbled crayfish (central box) in a German low-land lake. Organisms are not to scale.

804

805